# **APPENDIX: SUPPLEMENTAL TABLES**

The table numbers in this appendix are correlated with figure numbers throughout the document.

# Cancer in Rhode Island: Cancer Trends

| Table 5-     | Table 5-1. Cancer incidence by year and sex for all cancers combined   |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
|--------------|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Annual i     | Annual invasive* cancer incidence rates** for all cancers combined, RI and US, 1973-2000.  |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
| Male         |  |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
|              | 1973   | 1974  | 1975  | 1976  | 1977  | 1978  | 1979  | 1980  | 1981  | 1982  | 1983  | 1984  | 1985  | 1986  | 1987  |
| RI           | -  | -     | -     | -     | -     | -     | -     | -     | -     | -     | -     | -     | -     | -     | 566.4 |
| US           | 448.5  | 457.9 | 466.4 | 480.9 | 486.3 | 488.3 | 496.2 | 505.4 | 510.6 | 510.7 | 519.6 | 525.1 | 530.7 | 537.9 | 562.8 |
|              | 1988   | 1989  | 1990  | 1991  | 1992  | 1993  | 1994  | 1995  | 1996  | 1997  | 1998  | 1999  | 2000  | 2001  |       |
| RI           | 559.7  | 535.0 | 561.3 | 596.3 | 614.1 | 625.5 | 609.5 | 616.9 | 636.9 | 634.0 | 623.8 | 631.5 | 645.8 | 633.7 |       |
| US           | 557.9  | 566.5 | 592.0 | 636.5 | 656.8 | 620.6 | 587.9 | 568.7 | 568.2 | 570.5 | 566.0 | 569.1 | 560.2 | -     |       |
| Female       |  |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
|              | 1973   | 1974  | 1975  | 1976  | 1977  | 1978  | 1979  | 1980  | 1981  | 1982  | 1983  | 1984  | 1985  | 1986  | 1987  |
| RI           | -  | -     | -     | -     | -     | -     | -     | -     | -     | -     | -     | -     | -     | -     | 407.1 |
| US           | 349.5  | 370.5 | 365.8 | 367.2 | 363.4 | 362.1 | 365.3 | 367.6 | 376.8 | 375.9 | 380.4 | 392.0 | 402.8 | 402.0 | 413.2 |
|              | 1988   | 1989  | 1990  | 1991  | 1992  | 1993  | 1994  | 1995  | 1996  | 1997  | 1998  | 1999  | 2000  | 2001  |       |
| RI           | 419.8  | 411.3 | 413.4 | 425.6 | 433.8 | 432.6 | 442.0 | 446.3 | 450.6 | 458.5 | 446.5 | 470.3 | 454.2 | 428.6 |       |
| US           | 410.9  | 410.4 | 417.2 | 420.4 | 417.1 | 410.2 | 415.2 | 416.1 | 418.7 | 428.3 | 432.5 | 426.3 | 413.8 | -     |       |
| * ** Source: | Invasive includes the following stages of disease at diagnosis: local, regional, distant, and unknown.  Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.  urce: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER*Stat. |       |       |       |       |       |       |       |       |       |       |       |       |       |       |

| Table 5-2 | able 5-2. Cancer mortality by year and sex for all cancers combined   |       |       |       |           |       |       |       |       |       |       |       |       |       |       |       |
|-----------|---|-------|-------|-------|-----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
|           | Annual cancer mortality rates* for all cancers combined, RI and US, 1969-2000.                              |       |       |       |           |       |       |       |       |       |       |       |       |       |       |       |
| Male      | Male  |       |       |       |           |       |       |       |       |       |       |       |       |       |       |       |
|           | 1969  | 1970  | 1971  | 1972  | 1973      | 1974  | 1975  | 1976  | 1977  | 1978  | 1979  | 1980  | 1981  | 1982  | 1983  | 1984  |
| RI        | 288.6   | 303.1 | 286.3 | 280.1 | 287.<br>6 | 283.5 | 294.5 | 315.3 | 319.9 | 307.4 | 298.1 | 286.1 | 297.9 | 304   | 308.2 | 303.3 |
| US        | 247.6   | 249.3 | 251.7 | 255.4 | 255.0     | 257.9 | 258.4 | 262.2 | 264.2 | 267.1 | 268.7 | 271.1 | 269.8 | 272.9 | 274.1 | 275.1 |
|           | 1985  | 1986  | 1987  | 1988  | 1989      | 1990  | 1991  | 1992  | 1993  | 1994  | 1995  | 1996  | 1997  | 1998  | 1999  | 2000  |
| RI        | 295.4   | 296.6 | 283.0 | 299.6 | 311.3     | 295.6 | 290.3 | 292.6 | 274.8 | 284.8 | 285.2 | 284.0 | 274.5 | 281.5 | 260.7 | 255.6 |
| US        | 275.7   | 276.0 | 276.3 | 276.9 | 278.5     | 279.8 | 279.1 | 276.5 | 275.9 | 272.1 | 268.5 | 263.7 | 258.1 | 253.6 | 252.8 | 249.8 |
| Female    |   |       |       |       |           |       |       |       |       |       |       |       |       |       |       |       |
|           | 1969  | 1970  | 1971  | 1972  | 1973      | 1974  | 1975  | 1976  | 1977  | 1978  | 1979  | 1980  | 1981  | 1982  | 1983  | 1984  |
| RI        | 172.0   | 173.2 | 174.1 | 182.1 | 173.1     | 165.6 | 173.5 | 181.1 | 189.9 | 181.0 | 175.7 | 177.3 | 172.5 | 175.9 | 180.0 | 185.9 |
| US        | 163.2   | 163.0 | 162.2 | 162.7 | 160.8     | 161.1 | 160   | 162.7 | 162.9 | 163.8 | 163.1 | 165.8 | 165.9 | 167.4 | 168.3 | 170.5 |
|           | 1985  | 1986  | 1987  | 1988  | 1989      | 1990  | 1991  | 1992  | 1993  | 1994  | 1995  | 1996  | 1997  | 1998  | 1999  | 2000  |
| RI        | 184.9   | 182.5 | 177.5 | 186.1 | 179.4     | 183.9 | 177.5 | 187.0 | 180.0 | 176.5 | 182.7 | 183.5 | 181.5 | 173.7 | 181.2 | 177.4 |
| US        | 171.0   | 171.6 | 171.5 | 172.5 | 174.3     | 174.6 | 175.3 | 174.4 | 174.5 | 174.1 | 173.4 | 171.2 | 169.0 | 166.9 | 167.2 | 167.3 |
| * Source: | rates are age-adjusted to the year 2000 os standard population, expressed as deaths per 100,000 population. |       |       |       |           |       |       |       |       |       |       |       |       |       |       |       |

Cancer in Rhode Island

# **Cancer in Rhode Island: Cancer Disparities**

### Table 5-3. Cancer incidence by sex for all cancers combined

Cancer incidence rates\* by sex for all cancers combined, RI and US, 1996-2000.

| Sex  | RI    | US    |  |  |  |  |
|--|-------|-------|--|--|--|--|
| Male   | 627.0 | 566.7 |  |  |  |  |
| Female   | 452.2 | 423.9 |  |  |  |  |
| * Pates are age-adjusted to the year 2000 US standard population |       |       |  |  |  |  |

Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER\*Stat.

| Table 5-4. Cancer mortality by sex for all cancers combined                    |
|--|
| Cancer mortality rates* by sex for all cancers combined, RI and US, 1996-2000. |

| Sex  | RI    | US    |  |  |  |  |  |
|--|-------|-------|--|--|--|--|--|
| Male   | 270.7 | 255.5 |  |  |  |  |  |
| Female   | 179.0 | 168.3 |  |  |  |  |  |
| * Pates are ago adjusted to the year 2000 US standard population |       |       |  |  |  |  |  |

Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

Source: Office of Vital Records, HEALTH; SEER US Mortality 1969-2000 Data; calculated with SEER\*Stat.

### Table 5-5. Cancer incidence by age and sex for all cancers combined

Invasive\* cancer incidence rates\*\* by age and sex for all cancers combined, RI and US, 1996-2000.

|        | 00-<br>04 | 05-<br>09 | 10-<br>14 | 15-<br>19 | 20-<br>24 | 25-<br>29 | 30-<br>34 | 35-<br>39 | 40-<br>44 | 45-<br>49 | 50-<br>54 | 55-<br>59 | 60-<br>64 | 65-<br>69 | 70-<br>74 | 75-<br>79 | 80-<br>84 | 85+    |
|--------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|--------|
| Male   |           |           |           |           |           |           |           |           |           |           |           |           |           |           |           |           |           |        |
| RI     | 21.1      | 9.4       | 20.2      | 16.1      | 34.5      | 42.9      | 78.3      | 97.4      | 173.9     | 323.9     | 623.0     | 1067.6    | 1905.9    | 2636.3    | 3328.2    | 3596.2    | 3523.1    | 3727.2 |
| US     | 21.5      | 12.2      | 13.0      | 21.0      | 34.4      | 49.6      | 70.5      | 100.7     | 159.2     | 285.0     | 547.0     | 993.4     | 1630.5    | 2392.4    | 3046.7    | 3355.4    | 3495.3    | 3441.4 |
| Female |           |           |           |           |           |           |           |           |           |           |           |           |           |           |           |           |           |        |
| RI     | 24.2      | 18.8      | 13.3      | 19.8      | 37.4      | 51.8      | 110.6     | 201.4     | 338.0     | 527.5     | 750.9     | 994.2     | 1415.0    | 1586.5    | 1910.4    | 2116.9    | 2160.2    | 2140.0 |
| US     | 19.8      | 9.8       | 12.2      | 22.0      | 42.7      | 69.5      | 113.3     | 188.1     | 321.1     | 501.4     | 708.5     | 960.6     | 1225.3    | 1544.9    | 1843.0    | 2040.8    | 2138.5    | 1996.7 |

Invasive includes the following stages of disease at diagnosis: local, regional, distant, and unknown.
Rates are age-specific, expressed as cases per 100,000 population.

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER\*Stat.

### Table 5-6. Cancer mortality by age and sex for all cancers combined

Cancer mortality rates\* by age and sex for all cancers combined, RI and US, 1996-2000.

|        | 04  | 09  | 14  | 19  | 24  | 29  | 34   | 39   | 44   | 49    | 54    | 59    | 64    | 69    | 74     | 79     | 84     | 85+    |
|--------|---|-----|-----|-----|-----|-----|------|------|------|-------|-------|-------|-------|-------|--------|--------|--------|--------|
| Male   |   |     |     |     |     |     |      |      |      |       |       |       |       |       |        |        |        |        |
| RI     | 1.8   | 4.4 | 3.5 | 3.3 | 5.8 | 8.8 | 11.9 | 27.8 | 42.3 | 94.4  | 184.9 | 304.0 | 570.8 | 876.5 | 1328.0 | 1718.8 | 2220.3 | 3170.5 |
| US     | 2.7   | 2.7 | 2.8 | 4.2 | 6.1 | 7.8 | 12.5 | 22.4 | 46.3 | 95.1  | 180.8 | 334.0 | 561.1 | 863.5 | 1232.9 | 1601.3 | 2069.3 | 2716.5 |
| Female |   |     |     |     |     |     |      |      |      |       |       |       |       |       |        |        |        |        |
| RI     | 1.3   | 2.9 | 4.3 | 2.3 | 5.8 | 7.5 | 14.9 | 35.0 | 49.1 | 102.8 | 176.1 | 287.4 | 453.3 | 608.8 | 794.8  | 1022.4 | 1205.8 | 1593.3 |
| US     | 2.4   | 2.3 | 2.3 | 3.0 | 4.3 | 7.4 | 14.6 | 29.2 | 54.8 | 96.0  | 164.9 | 267.2 | 408.1 | 572.5 | 773.8  | 956.6  | 1163.9 | 1423.9 |
| *      | Rates are age-specific, expressed as deaths per 100,000 population. |     |     |     |     |     |      |      |      |       |       |       |       |       |        |        |        |        |

00- 05- 10- 15- 20- 25- 30- 35- 40- 45- 50- 55- 60- 65- 70- 75- 80-

Source: SEER US Mortality 1969-2000 Data; SEER web query system.

### Table 5-7. Cancer incidence by race and sex for all cancers combined

Cancer incidence rates\* by race and sex for all cancers combined, RI and US, 1987-2000.

|    | M     | ale   | Fer   | Female |  |  |  |
|----|-------|-------|-------|--------|--|--|--|
|    | White | Black | White | Black  |  |  |  |
| RI | 631.5 | 659.1 | 483.4 | 438.6  |  |  |  |
| US | 604.1 | 731.9 | 501.5 | 460.1  |  |  |  |

\* Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER\*Stat.

### Table 5-8. Cancer mortality by race and sex for all cancers combined

Cancer mortality rates\* by race and sex for all cancers combined, RI and US, 1987-2000.

|    | M     | ale   | Female |       |  |  |  |
|----|-------|-------|--------|-------|--|--|--|
|    | White | Black | White  | Black |  |  |  |
| RI | 285.8 | 380.3 | 183.7  | 234.3 |  |  |  |
| US | 261.0 | 376.3 | 170.2  | 201.8 |  |  |  |

Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

Source: Office of Vital Records, HEALTH; SEER US Mortality 1969-2000 Data; calculated with SEER\*Stat.

### **Breast Cancer**

### Table 6-1. Female breast cancer incidence by year

Average annual invasive\* breast cancer incidence rates\*\* among females, RI and US, 1987-2001\*\*\*.

|    | 1989  | 1990  | 1991  | 1992  | 1993  | 1994  | 1995  | 1996  | 1997  | 1998   | 1999  |
|----|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|-------|
| RI | 132.6 | 131.5 | 130.1 | 128.8 | 129.9 | 130.6 | 130.8 | 131.2 | 131.5 | 131.1  | 131.2 |
| US | 131.5 | 131.1 | 130.7 | 131.4 | 131.6 | 131.6 | 132.7 | 135.0 | 136.6 | 137.1* | NA    |

Invasive includes the following stages of disease at diagnosis: local, regional, distant, and unknown.

\*\* Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

Source: RICR, HEALTH – calculated with SEER\*Stat; SEER Cancer Statistics Review, 1973-1999; 1998 US data is from SEER Public-Use 1973-2000 Data – calculated with SEER\*Stat.

### Table 6-2. Female breast cancer mortality by year

Average annual breast cancer mortality rates\* among females, RI and US, 1987-2000\*\*

|    | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998  |
|----|------|------|------|------|------|------|------|------|------|-------|
| RI | 37.2 | 37.9 | 36.5 | 35.6 | 35.2 | 34.2 | 32.2 | 32.2 | 31.1 | 29.2  |
| US | 33.0 | 32.9 | 32.5 | 32.1 | 31.6 | 31.0 | 30.4 | 29.6 | 28.8 | 27.7* |

Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

\*\* Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

Source: CDC WONDER, CDC;

1998 US data is from SEER US Mortality 1969-2000 Data - calculated with SEER\*Stat.

### Table 6-3. Female breast cancer incidence by race

Average annual invasive breast cancer incidence rates\* by race among females, RI and US, 1987-2000.

|    | White | Black |
|----|-------|-------|
| RI | 156.3 | 129.0 |
| US | 162.9 | 139.9 |

\* Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER\*Stat.

### Table 6-4. Female breast cancer mortality by race

Average annual breast cancer mortality rates\* by race among females, RI and US, 1987-2000.

|    | White                     | Black                       |
|----|---------------------------|-----------------------------|
| RI | 34.1                      | 40.5                        |
| US | 30.1                      | 36.9                        |
| *  | Rates are age-adjusted to | o the year 2000 US standard |

population, expressed as deaths per 100,000 population. Source: RICR, HEALTH; SEER US Mortality 1969-2000 data; calculated

with SEER\*Stat.

### Table 6-6. Female breast cancer screening by year

Percent of female respondents, age 40 and older, who report that they have had a mammogram within the past two years, RI and US, 1990-2001.

|    | 1990                          | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |  |
|----|-------------------------------|------|------|------|------|------|------|------|------|------|------|------|--|
| RI | 71                            | 71   | 66   | 73   | NA   | 72   | 76   | 81   | 80   | 83   | 84   | 85   |  |
| US | 58                            | 62   | 63   | 66   | 67   | 69   | 69   | 70   | 72   | 73   | 76   | NA   |  |
| •  | DI DEFON LIFALTILI DEFON OD O |      |      |      |      |      |      |      |      |      |      |      |  |

Source: RI-BRFSS, HEALTH; BRFSS, CDC.

### Table 6-7. Female breast cancer cases in ACOS programs by year

Percent of breast cancer cases that were or are treated in ACOS-approved cancer treatment programs by year among females, RI, 1989-2000.

| 1989                  | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|-----------------------|------|------|------|------|------|------|------|------|------|------|------|
| 52                    | 53   | 55   | 70   | 70   | 70   | 69   | 69   | 66   | 81   | 81   | 89   |
| Source: RICR, HEALTH. |      |      |      |      |      |      |      |      |      |      |      |

### Table 6-8. Female breast cancer cases with AJCC staging by year

Percent of breast cancer cases staged with AJCC staging methodology by year among females, RI, 1989-2000.

| 1989      | 1990                  | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|-----------|-----------------------|------|------|------|------|------|------|------|------|------|------|
| 73        | 74                    | 71   | 75   | 90   | 93   | 93   | 94   | 92   | 92   | 92   | 95   |
| Source: I | Source: RICR, HEALTH. |      |      |      |      |      |      |      |      |      |      |

### Table 6-9. Female breast cancer survival rates by race and stage

Five-year relative invasive breast cancer survival rate\*s by race and stage of disease at diagnosis among females, US, 1992-1999.

|       | In situ**   | Localized | Regional | Distant | Unstaged |  |  |  |  |  |  |  |
|-------|---|-----------|----------|---------|----------|--|--|--|--|--|--|--|
| White | 100.0   | 97.6      | 80.3     | 24.6    | 56.3     |  |  |  |  |  |  |  |
| Black | 100.0   | 89.7      | 66.0     | 15.2    | 52.0     |  |  |  |  |  |  |  |
| *     | * Survival rates are relative rates expressed as percents |           |          |         |          |  |  |  |  |  |  |  |

In situ cases are not considered invasive. Source: SEER Cancer Statistics Review, 1975-2000.

### **Cervical Cancer**

### Table 7-1. Cervical cancer incidence by year

Average annual invasive\* cervical cancer incidence rates\*\* among females, RI and US, 1987-2001\*\*\*.

|    | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|----|------|------|------|------|------|------|------|------|------|------|------|
| RI | 10.0 | 10.8 | 11.0 | 11.2 | 11.6 | 11.8 | 11.4 | 10.7 | 10.4 | 9.7  | 8.8  |
| US | 10.4 | 10.4 | 10.2 | 10.0 | 9.6  | 9.5  | 9.4  | 9.3  | 9.0  | 8.7* | NA   |

- Invasive includes the following stages of disease at diagnosis: local, regional, distant, and unknown.
- Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.
- Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

RICR, HEALTH - calculated with SEER\*Stat; SEER Cancer Statistics Review, 1973-1999;

1998 US data is from SEER Public-Use 1973-2000 Data - calculated with SEER\*Stat.

### Table 7-2. Cervical cancer mortality by year

Average annual cervical cancer mortality rates\* among females, RI and US, 1987-2000\*\*.

|    | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
|----|------|------|------|------|------|------|------|------|------|------|
| RI | 2.6  | 2.8  | 3.0  | 3.0  | 2.8  | 3.3  | 3.2  | 2.9  | 2.9  | 2.8  |
| US | 3.6  | 3.6  | 3.5  | 3.5  | 3.4  | 3.4  | 3.3  | 3.2  | 3.1  | 3.0* |

- Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.
- Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

Source: CDC WONDER, CDC;

1998 US data is from SEER US Mortality 1969-2000 Data - calculated with SEER\*Stat.

### Table 7-3. Cervical cancer incidence by race

Average annual invasive cervical cancer incidence rates\* by race among females, RI and US, 1987-2000.

|    | White | Black |
|----|-------|-------|
| RI | 10.2  | 18.1  |
| US | 8.0   | 12.7  |
| 4  |       |       |

Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated

with SEER\*Stat.

### Table 7-4. Cervical cancer mortality by race

Average annual cervical cancer mortality rates\* by race among females, RI and US, 1987-2000.

|    | White | Black |
|----|-------|-------|
| RI | 2.9   | 6.4   |
| US | 2.9   | 7.0   |

\* Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

Source: RICR, HEALTH; SEER US Mortality 1969-2000 data; calculated with SEER\*Stat.

### Table 7-5. Cervical cancer screening by year

Percent of female respondents, with uterine cervix, age 18 and older, who report that they have had a pap smear within the last three years, RI and US, 1992-2001.

|        | 1992                             | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |  |  |
|--------|----------------------------------|------|------|------|------|------|------|------|------|------|--|--|
| RI     | 80.0                             | 83.9 | NA   | 82.9 | 82.9 | 87.4 | 85.2 | 86.0 | 88.5 | 88.2 |  |  |
| US     | 84.1                             | 84.7 | 85.4 | 84.5 | 84.6 | 84.6 | 84.8 | 85.5 | 86.8 | NA   |  |  |
| Couroo | Course. DI DDCC LIFALTH DDCC CDC |      |      |      |      |      |      |      |      |      |  |  |

Source: RI-BRFSS, HEALTH; BRFSS, CDC.

### Table 7-6. Cervical cancer cases in ACOS programs by year

Percent of cervical cancer cases that were or are treated in ACOS-approved cancer treatment programs by year among females, RI, 1989-2000.

| 1989      | 1990       | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|-----------|------------|------|------|------|------|------|------|------|------|------|------|
| 45        | 33         | 35   | 39   | 44   | 41   | 36   | 41   | 32   | 91   | 98   | 97   |
| Source: I | RICR, HEAL | .TH. |      |      |      |      |      |      |      |      |      |

### Table 7-7. Cervical cancer cases with AJCC staging by year

Percent of cervical cancer cases staged with AJCC staging methodology by year among females, RI, 1989-2000.

| i ciccin o | referred by the real career cases staged with 1000 staging methodology by year among termines, Nr, 1707 2000. |      |      |      |      |      |      |      |      |      |      |  |  |  |
|------------|---|------|------|------|------|------|------|------|------|------|------|--|--|--|
| 1989       | 1990  | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |  |  |  |
| 72         | 77  | 81   | 73   | 84   | 97   | 90   | 93   | 94   | 87   | 92   | 87   |  |  |  |
| Source:    | Source: RICR, HEALTH.   |      |      |      |      |      |      |      |      |      |      |  |  |  |

### Table 7-8. Cervical cancer survival rates by race and stage

Five-year relative invasive cervical cancer survival rates by race and stage of disease at diagnosis among females, US, 1992-1999.

|       | Localized  | Regional | Distant | Unstaged |  |  |  |  |  |
|-------|--|----------|---------|----------|--|--|--|--|--|
| White | 92.6   | 51.8     | 17.5    | 59.0     |  |  |  |  |  |
| Black | 87.2   | 41.4     | 11.6    | 50.9     |  |  |  |  |  |
| *     | Survival rates are relative rates expressed as percents. |          |         |          |  |  |  |  |  |

Source: SEER Cancer Statistics Review, 1975-2000.

# **Colorectal Cancer**

| Tables 8        | -1 and 8-2.            | Colorecta             | l cancer ir               | ncidence k              | y year an   | d sex                     |                           |                        |             |                      |         |
|-----------------|------------------------|-----------------------|---------------------------|-------------------------|---|---------------------------|---------------------------|------------------------|-------------|----------------------|---------|
| Average         | e annual in            | vasive* col           | lorectal ca               | ncer incia              | lence rate.   | s** by sex,               | RI and US,                | 1987-2001'             | ***         |                      |         |
|                 | 1989                   | 1990                  | 1991                      | 1992                    | 1993  | 1994                      | 1995                      | 1996                   | 1997        | 1998                 | 1999    |
| Male            |                        |                       |                           |                         |   |                           |                           |                        |             |                      |         |
| RI              | 94.1                   | 91.0                  | 85.8                      | 84.4                    | 80.9  | 78.3                      | 77.8                      | 77.5                   | 75.7        | 75.9                 | 76.4    |
| US              | 75.0                   | 73.6                  | 72.2                      | 70.4                    | 68.2  | 66.6                      | 66.1                      | 65.6                   | 65.1        | 65.4*                | NA      |
| Female          |                        |                       |                           |                         |   |                           |                           |                        |             |                      |         |
| RI              | 58.9                   | 57.6                  | 59.0                      | 57.9                    | 54.8  | 54.5                      | 56.5                      | 55.3                   | 56.6        | 56.2                 | 55.4    |
| US              | 51.7                   | 51.0                  | 50.3                      | 49.4                    | 48.5  | 47.8                      | 47.6                      | 47.7                   | 47.6        | 47.4*                | NA      |
| * ** ** Source: | Rates are<br>Rates are | age-adjus<br>based on | ted to the<br>five years' | year 2000<br>data (e.g. | disease at<br>US standa<br>., 1989 = 19<br>; SEER Can | rd populat<br>187-1991; 1 | ion, expres<br>998 = 1996 | ssed as ca:<br>-2000). | ses per 100 | known.<br>),000 popu | lation. |

| Tables 8     | -3 and 8-4. (  | Colorectal c | ancer mort  | ality by yea | ar and sex  |              |        |      |      |       |
|--------------|--|--------------|-------------|--------------|-------------|--------------|--------|------|------|-------|
| Average      | e annual col   | orectal car  | cer mortali | ty rates* by | sex, RI and | US, 1987-200 | 00***. |      |      |       |
|              | 1989   | 1990         | 1991        | 1992         | 1993        | 1994         | 1995   | 1996 | 1997 | 1998  |
| Male         | •  |              |             |              |             |              |        |      |      |       |
| RI           | 37.0   | 36.9         | 34.5        | 34.1         | 32.7        | 33.1         | 31.4   | 31.9 | 30.2 | 28.9  |
| US           | 30.3   | 29.9         | 29.4        | 29.0         | 28.4        | 27.9         | 27.3   | 26.7 | 26.1 | 25.8* |
| Female       |  |              |             |              |             |              |        |      |      |       |
| RI           | 22.7   | 22.7         | 22.2        | 21.0         | 20.7        | 20.6         | 20.2   | 19.5 | 20.2 | 20.1  |
| US           | 20.9   | 20.5         | 20.3        | 20.0         | 19.7        | 19.4         | 19.1   | 18.8 | 18.5 | 18.0* |
| * ** Source: | Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.  Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).  CDC WONDER, CDC;  1998 US data is from SEER US Mortality 1969-2000 Data – calculated with SEER*Stat. |              |             |              |             |              |        |      |      |       |

| Table 8-            | 5. Colorectal cancer i   | incidence by race a   | nd sex                |                     |  |  |  |  |  |  |  |  |
|---------------------|--|-----------------------|-----------------------|---------------------|--|--|--|--|--|--|--|--|
| Average<br>1987-200 | e annual invasive colo<br>00.  | orectal cancer incide | ence rates* by race a | and sex, RI and US, |  |  |  |  |  |  |  |  |
|                     | Male Female  |                       |                       |                     |  |  |  |  |  |  |  |  |
|                     | White  | Black                 | White                 | Black               |  |  |  |  |  |  |  |  |
| RI                  | 90.9   | 73.9                  | 61.3                  | 60.7                |  |  |  |  |  |  |  |  |
| US                  | 73.8   | 79.1                  | 51.5                  | 59.9                |  |  |  |  |  |  |  |  |
| * Source:           | Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.  RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER*Stat. |                       |                       |                     |  |  |  |  |  |  |  |  |

### Table 8-6. Colorectal cancer mortality by race and sex

Average annual colorectal cancer mortality rates\* by race and sex, RI and US, 1987-2000.

|    | M     | ale   | Female |       |  |
|----|-------|-------|--------|-------|--|
|    | White | Black | White  | Black |  |
| RI | 34.0  | 26.9  | 21.8   | 29.0  |  |
| US | 27.8  | 35.4  | 19.0   | 25.4  |  |

\* Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

Source: RICR, HEALTH; SEER US Mortality 1969-2000 data; calculated with SEER\*Stat.

### Tables 8-9 and 8-10. Colorectal cancer screening by year and sex

Percent of respondents 40 and older who have ever been screened for colorectal cancer (proctoscopic exam, sigmoidoscopy, or colonoscopy), by sex, RI and US, 1995-1999.

| VI      |                   | 1.57        |      | ,    |      |
|---------|-------------------|-------------|------|------|------|
|         | 1995              | 1996        | 1997 | 1998 | 1999 |
| Male    |                   |             |      |      |      |
| RI      | 32.0              | NA          | 38.2 | NA   | 37.4 |
| US      | 32.1              | NA          | 34.6 | NA   | 33.9 |
| Female  |                   |             |      |      |      |
| RI      | 25.6              | NA          | 31.6 | NA   | 37.2 |
| US      | 27.1              | NA          | 30.4 | NA   | 32.6 |
| Source: | RI-BRFSS, HEALTH; | BRFSS, CDC. |      |      |      |

### Table 8-11. Colorectal cancer cases in ACOS programs by year and sex

Percent of colorectal cancer cases that were or are treated in ACOS-approved cancer treatment programs by year and sex, RI, 1989-2000.

|         | 1989                 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|---------|----------------------|------|------|------|------|------|------|------|------|------|------|------|
| Male    | 49                   | 49   | 51   | 69   | 73   | 70   | 72   | 75   | 71   | 75   | 67   | 82   |
| Female  | 53                   | 47   | 50   | 69   | 69   | 67   | 70   | 71   | 70   | 72   | 70   | 83   |
| Source: | ource: RICR, HEALTH. |      |      |      |      |      |      |      |      |      |      |      |

### Table 8-12. Colorectal cancer cases with AJCC staging by year and sex

Percent of colorectal cancer cases staged with AJCC staging methodology by year and sex, RI, 1989-2000.

| I CICCIII ( | or colorec | nai carice | or Cases si | agea viii | 17150031 | aging me | unoaciog | iy by year | aria scx, | 111, 1707 21 | 500. |      |
|-------------|------------|------------|-------------|-----------|----------|----------|----------|------------|-----------|--------------|------|------|
|             | 1989       | 1990       | 1991        | 1992      | 1993     | 1994     | 1995     | 1996       | 1997      | 1998         | 1999 | 2000 |
| Male        | 65         | 64         | 68          | 68        | 89       | 92       | 92       | 92         | 91        | 95           | 90   | 92   |
| Female      | 65         | 59         | 62          | 68        | 87       | 90       | 96       | 90         | 90        | 90           | 92   | 90   |
| Sourco      | DICD HE    | \ I TLI    |             |           |          |          |          |            |           |              |      |      |

### Table 8-13. Male colorectal cancer survival rates by race and stage

Five-year relative invasive colorectal cancer survival rates by stage of disease at diagnosis among males, US, 1992-1999.

|         | Localized             | Regional  | Distant | Unstaged |  |  |  |  |  |
|---------|-----------------------|---|---------|----------|--|--|--|--|--|
| White   | 91.1                  | 65.7  | 9.0     | 39.7     |  |  |  |  |  |
| Black   | 83.0                  | 59.1  | 7.3     | 42.8     |  |  |  |  |  |
| *       | Survival rates are re | urvival rates are relative rates expressed as percents. |         |          |  |  |  |  |  |
| Source: | SEER Cancer Statis    | SEER Cancer Statistics Review, 1975-2000.               |         |          |  |  |  |  |  |

### Table 8-14. Female colorectal cancer survival rates by race and stage

Five-year relative invasive colorectal cancer survival rates by race and stage of disease at diagnosis among females, US, 1992-1999.

|       | Localized  | Regional | Distant | Unstaged |  |  |  |  |  |
|-------|--|----------|---------|----------|--|--|--|--|--|
| White | 90.0   | 66.9     | 9.6     | 31.6     |  |  |  |  |  |
| Black | 83.8   | 58.1     | 9.0     | 28.6     |  |  |  |  |  |
| *     | * Survival rates are relative rates expressed as percents. |          |         |          |  |  |  |  |  |

Source: SEER Cancer Statistics Review, 1975-2000.

# **Lung Cancer**

| Tables 9 | -1 and 9-2.  | Lung cand   | er inciden  | ce by yea   | r and sex   |             |              |              |             |            |        |
|----------|--|-------------|-------------|-------------|-------------|-------------|--------------|--------------|-------------|------------|--------|
|          | verage annual invasive* lung cancer incidence rates** by sex, RI and US, 1987-2001***. |             |             |             |             |             |              |              |             |            |        |
|          | 1989   | 1990        | 1991        | 1992        | 1993        | 1994        | 1995         | 1996         | 1997        | 1998       | 1999   |
| Male     |  |             |             |             |             |             |              |              |             |            |        |
| RI       | 105.1  | 106.2       | 107.3       | 106.4       | 106.6       | 110.7       | 109.2        | 107.6        | 106.8       | 104.6      | 102.3  |
| US       | 98.7   | 97.9        | 96.9        | 95.4        | 93.8        | 91.8        | 89.4         | 88.0         | 86.0        | 85.0*      | NA     |
| Female   |  |             |             |             |             |             |              |              |             |            |        |
| RI       | 46.3   | 49.6        | 50.2        | 52.4        | 54.4        | 57.0        | 57.9         | 60.6         | 61.9        | 62.5       | 61.3   |
| US       | 46.8   | 48.0        | 48.5        | 49.4        | 49.9        | 50.3        | 50.8         | 51.4         | 51.4        | 51.6*      | NA     |
| *        | Invasive in  | ncludes the | following   | stages of o | disease at  | diagnosis:  | local, regio | onal, distar | nt, and unl | nown.      |        |
| **       | Rates are  | age-adjus   | ted to the  | year 2000   | US standar  | d populat   | ion, expres  | sed as cas   | ses per 100 | ,000 popul | ation. |
| ***      | Rates are  | based on    | five years' | data (e.g.  | , 1989 = 19 | 87-1991; 1  | 998 = 1996   | -2000).      | -           | • •        |        |
| Source:  |  |             | ulated with | ٠ ٠         |             |             |              | ,            | );          |            |        |
|          | 1998 US da   | ata is from | SEER Public | c-Use 1973  | -2000 Data  | a – calcula | ted with SI  | EER*Stat.    |             |            |        |

| Tables 9     | -3 and 9-4. l  | ung cance  | r mortality k | by year and   | sex          |           |      |      |      |       |
|--------------|--|------------|---------------|---------------|--------------|-----------|------|------|------|-------|
| Average      | e annual lun   | g cancer m | ortality rate | es* by sex, R | l and US, 19 | 87-2000** |      |      |      |       |
|              | 1989   | 1990       | 1991          | 1992          | 1993         | 1994      | 1995 | 1996 | 1997 | 1998  |
| Male         |  |            |               |               |              |           |      |      |      |       |
| RI           | 91.9   | 92.0       | 92.7          | 91.6          | 92.3         | 91.5      | 91.3 | 90.7 | 89.5 | 85.9  |
| US           | 90.2   | 89.8       | 89.4          | 88.7          | 87.4         | 85.9      | 84.6 | 83.0 | 81.2 | 79.5* |
| Female       |  |            |               |               |              |           |      |      |      |       |
| RI           | 35.5   | 37.0       | 38.3          | 38.9          | 40.7         | 43.3      | 44.9 | 44.9 | 46.2 | 45.0  |
| US           | 35.6   | 36.8       | 37.9          | 38.7          | 39.4         | 40.0      | 40.5 | 40.9 | 41.1 | 40.7* |
| * ** Source: | Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.  Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).  CDC WONDER, CDC;  1998 US data is from SEER US Mortality 1969-2000 Data – calculated with SEER*Stat. |            |               |               |              |           |      |      |      |       |

### Table 9-5. Lung cancer incidence by race and sex

Average annual invasive lung cancer incidence rates\* by race and sex, RI and US, 1987-2000.

|    | M           | ale   | Female |       |  |  |  |
|----|-------------|-------|--------|-------|--|--|--|
|    | White Black |       | White  | Black |  |  |  |
| RI | 108.1       | 127.1 | 56.1   | 59.2  |  |  |  |
| US | 90.6        | 134.2 | 51.0   | 53.3  |  |  |  |

Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER\*Stat.

### Table 9-6. Lung cancer mortality by race and sex

Average annual lung cancer mortality rates\* by race and sex, RI and US, 1987-2000.

|    | M     | ale         | Female |       |  |  |  |
|----|-------|-------------|--------|-------|--|--|--|
|    | White | White Black |        | Black |  |  |  |
| RI | 90.3  | 117.7       | 41.5   | 52.5  |  |  |  |
| US | 83.2  | 115.8       | 39.2   | 37.9  |  |  |  |

Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

Source: RICR, HEALTH; SEER US Mortality 1969-2000 data; calculated with SEER\*Stat.

### Tables 9-9 and 9-10. Current smokers by year and sex

Percent of respondents 18 and older who have ever smoked 100 cigarettes in their lifetime and reported smoking every day or some days, by year and sex, RI and US, 1990-2001.

| aay o. o. | orric days, | , 25 5 0 4 | arra 00x, r |      | .,,,,,,, | •    |      |      |      |      |      |      |
|-----------|-------------|------------|-------------|------|----------|------|------|------|------|------|------|------|
|           | 1990        | 1991       | 1992        | 1993 | 1994     | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
| Male      |             |            |             |      |          |      |      |      |      |      |      |      |
| RI        | 27.3        | 24.9       | 24.2        | 25.9 | NA       | 24.0 | 25.6 | 25.5 | 24.0 | 23.1 | 23.8 | 25.8 |
| US        | 24.9        | 25.1       | 24.2        | 24.0 | 23.9     | 24.8 | 25.5 | 25.4 | 25.3 | 24.2 | 24.4 | 25.4 |
| Female    |             |            |             |      |          |      |      |      |      |      |      |      |
| RI        | 24.0        | 25.1       | 20.4        | 21.1 | NA       | 25.4 | 19.7 | 23.2 | 21.3 | 21.5 | 23.0 | 22.1 |
| US        | 21.3        | 21.3       | 21.0        | 21.1 | 21.6     | 20.9 | 21.9 | 21.1 | 20.9 | 20.8 | 21.2 | 21.2 |
| Source:   | RI-BRFSS,   | HEALTH; E  | BRFSS, CD   | C.   |          |      |      |      |      |      |      |      |

### Table 9-11. Lung cancer cases in ACOS programs by year and sex

Percent of lung cancer cases that were or are treated in ACOS-approved cancer treatment programs by year and sex, RI, 1989-2000.

|         | 1989                 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|---------|----------------------|------|------|------|------|------|------|------|------|------|------|------|
| Male    | 64                   | 64   | 60   | 75   | 78   | 77   | 80   | 76   | 77   | 77   | 80   | 88   |
| Female  | 53                   | 58   | 56   | 79   | 78   | 78   | 77   | 80   | 78   | 79   | 82   | 91   |
| Source: | purce: RICR, HEALTH. |      |      |      |      |      |      |      |      |      |      |      |

#### Table 9-12. Lung cancer cases with AJCC staging by year and sex Percent of lung cancer cases staged with AJCC staging methodology by year and sex, RI, 1989-2000. Male Female Source: RICR, HEALTH.

# Table 9-13. Male lung cancer survival rates by race and stage Invasive lung cancer five-year relative survival rates by stage of disease at diagnosis among males, US, 1992-1999. Localized Regional Distant Unstaged

|         | Localizoa         | Regional            | Distant        | onstaged |
|---------|-------------------|---------------------|----------------|----------|
| White   | 45.0              | 15.2                | 1.8            | 7.3      |
| Black   | 40.0              | 12.7                | 1.6            | 9.7      |
| *       | Survival rates ar | e relative rates ex | xpressed as pe | rcents.  |
| Source: | SEER Cancer Sta   | atistics Review, 19 | 775-2000.      |          |

| Table 9-1 | 14. Female lung o | cancer survival ra                      | ates by race an | ıd stage         |
|-----------|-------------------|---|-----------------|------------------|
|           |                   | e lung cancer sur<br>ong females, US, 1 |                 | ace and stage of |
|           | Localized         | Regional                                | Distant         | Unstaged         |
| White     | 53.6              | 17.7                                    | 2.2             | 9.4              |
| Black     | 47.7              | 15.0                                    | 2.4             | 8.7              |

\* Survival rates are relative rates expressed as percents.
Source: SEER Cancer Statistics Review, 1975-2000.

## Melanoma of Skin

| Tables 1 | 0-1 and 10  | -2. Melano  | ma of skin | incidence    | by year a  | nd sex     |             |           |            |      |       |
|----------|-------------|-------------|------------|--------------|------------|------------|-------------|-----------|------------|------|-------|
| Averag   | e annual in | vasive* me  | elanoma o  | f skinincide | ence rates | ** by year | and sex, Ri | and US, 1 | 987-2001** | *    |       |
|          | 1989        | 1990        | 1991       | 1992         | 1993       | 1994       | 1995        | 1996      | 1997       | 1998 | 1999  |
| Male     |             |             |            |              |            |            |             |           |            |      |       |
| RI       | 15.4        | 16.3        | 17.4       | 17.2         | 18.5       | 19.9       | 19.6        | 20.5      | 22.4       | 22.7 | 23.8  |
| US       | 21.9        | 23.0        | 24.1       | 25.4         | 26.8       | 28.4       | 29.8        | 31.7      | 33.1       | 34.5 | NA    |
| Female   |             |             |            |              |            |            |             |           |            |      |       |
| RI       | 8.9         | 10.4        | 10.4       | 11.4         | 12.6       | 12.9       | 12.7        | 12.9      | 12.9       | 13.1 | 13.8  |
| US       | 15.7        | 16.1        | 16.5       | 17.1         | 17.9       | 18.7       | 19.7        | 20.9      | 21.9       | 22.6 | NA    |
| *        |             | ncludes the |            |              |            |            |             |           |            |      | ation |

Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER\*Stat.

#### Tables 10-3 and 10-4. Melanoma of skin mortality by year and sex Average annual melanoma of skin mortality rates\* by year and sex, RI and US, 1987-2000\*\*. 1990 1989 1991 1992 1993 1994 1995 1996 1997 1998 Male RΙ 4.0 4.0 4.0 3.6 3.3 3.6 3.4 3.4 3.6 3.3 US 3.8 3.8 3.8 3.8 3.9 3.9 3.9 4.0 4.0 3.9 Female 1.5 1.7 1.9 1.7 2.0 2.5 RΙ 1.7 1.6 2.0 2.2 US 1.9 1.9 1.9 1.9 1.8 1.8 1.8 1.8 1.8 1.8

### Table 10-5. Melanoma of skin incidence by race and sex

Average annual invasive melanoma of skin incidence rates\* by race and sex, RI and US, 1987-2000.

|    | M     | ale   | Female |       |  |  |  |
|----|-------|-------|--------|-------|--|--|--|
|    | White | Black | White  | Black |  |  |  |
| RI | 25.7  | 0.3   | 16.6   | 0.5   |  |  |  |
| US | 32.6  | 1.8   | 22.2   | 1.3   |  |  |  |

Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER\*Stat.

### Table 10-6. Melanoma of skin mortality by race and sex

Average annual melanoma of skin mortality rates\* by race and sex, RI and US, 1987-2000.

|    | M     | ale         | Female |       |  |  |  |
|----|-------|-------------|--------|-------|--|--|--|
|    | White | White Black |        | Black |  |  |  |
| RI | 3.8   | 1.5         | 2.0    | 1.0   |  |  |  |
| US | 4.3   | 0.5         | 2.1    | 0.5   |  |  |  |

Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

Source: RICR, HEALTH; SEER US Mortality 1969-2000 data; calculated with SEER\*Stat.

### Table 10-7. Melanoma of skin cases in ACOS programs by year and sex

Percent of melanoma of skin cases that were or are treated in ACOS-approved cancer treatment programs by year and sex, RI, 1989-2000.

|         | 1989     | 1990  | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|---------|----------|-------|------|------|------|------|------|------|------|------|------|------|
| Male    | 57       | 67    | 56   | 87   | 86   | 87   | 82   | 85   | 84   | 84   | 86   | 93   |
| Female  | 55       | 71    | 72   | 85   | 78   | 82   | 79   | 88   | 84   | 86   | 88   | 92   |
| Source: | RICR, HE | ALTH. |      |      |      |      |      |      |      |      |      |      |

<sup>\*</sup> Rates are age-adjusted to the year 2000 US standard population, expressed as d eaths per 100,000 population.

\*\* Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

Source: Office of Vital Records, HEALTH; SEER US Mortality 1969-2000 Data; calculated with SEER\*Stat.

#### Table 10-8. Melanoma of skin cases with AJCC staging by year and sex Percent of melanoma of skin cases staged with AJCC staging methodology by year and sex, RI, 1989-2000. 1989 1990 1991 1992 1993 1994 1995 1996 1997 1999 2000 Male 53 50 56 37 42 66 67 64 68 43 28 Female 60 50 56 58 55 33 35 Source: RICR, HEALTH.

#### Table 10-9. Male melanoma of skin survival rates by race and stage Five-year relative invasive melanoma of skin survival rates\* by race and stage of disease at diagnosis among males, US, 1992-1999. Localized Regional Distant Unstaged White 56.0 12.4 76.2 96.5 Black 85.8\*\* NA NA NA Survival rates are relative rates expressed as percents. The standard error of the survival rate is greater than 10 percentage points. NΑ Statistic could not be calculated

Source: SEER Cancer Statistics Review, 1975-2000.

| Table 10                            | -10. Female mel  | anoma of skin sur                       | vival rates by r | ace and stage |  |  |  |  |  |  |  |
|-------------------------------------|--|---|------------------|---------------|--|--|--|--|--|--|--|
| _                                   | Five-year relative invasive melanoma of skin survival rates* by race and stage of disease at diagnosis among females, US, 1992-1999. |   |                  |               |  |  |  |  |  |  |  |
| Localized Regional Distant Unstaged |  |   |                  |               |  |  |  |  |  |  |  |
| White                               | 97.1 67.1 16.4 85.0  |   |                  |               |  |  |  |  |  |  |  |
| Black                               | 84.9**   | NA                                      | NA               | NA            |  |  |  |  |  |  |  |
| *                                   | survival rates are relative rates expressed as percents.   |   |                  |               |  |  |  |  |  |  |  |
| NA<br>Source:                       |  | ot be calculated<br>atistics Review, 19 |                  |               |  |  |  |  |  |  |  |

# **Oropharyngeal Cancer**

| Tables 1                                | 1-1 and 11-   | -2. Oropha  | ryngeal ca | ancer incid | lence by y | ear and se | ex          |              |          |      |      |  |
|---|---|---|------------|-------------|------------|------------|-------------|--------------|----------|------|------|--|
| Averag                                  | e annual in   | vasive* orc   | pharynge   | al cancer   | incidence  | rates** by | sex, RI and | I US, 1987-2 | 2001***. |      |      |  |
|   | 1989  | 1990  | 1991       | 1992        | 1993       | 1994       | 1995        | 1996         | 1997     | 1998 | 1999 |  |
| Male                                    |   |   |            |             |            |            |             |              |          |      |      |  |
| RI                                      | 17.6  | 16.6  | 14.9       | 14.7        | 14.8       | 15.0       | 15.6        | 16.2         | 16.8     | 16.7 | 15.1 |  |
| US                                      | 20.1  | 19.7  | 19.8       | 19.6        | 19.2       | 18.9       | 18.6        | 18.1         | 17.6     | 17.2 | NA   |  |
| Female                                  | 1   |   |            |             |            |            |             |              |          |      |      |  |
| RI                                      | 6.6   | 7.0   | 6.8        | 7.5         | 7.3        | 6.7        | 6.9         | 6.5          | 6.5      | 6.3  | 6.5  |  |
| US                                      | 7.5   | 7.4   | 7.4        | 7.4         | 7.4        | 7.3        | 7.3         | 7.2          | 7.1      | 6.9  | NA   |  |
| * | Invasive includes the following stages of disease at diagnosis: local, regional, distant, and unknown. Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population. |   |            |             |            |            |             |              |          |      |      |  |
| Source:                                 |   | Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).  RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER*Stat. |            |             |            |            |             |              |          |      |      |  |

| Tables 11 | ables 11-3 and 11-4. Oropharyngeal cancer mortality by year and sex                  |      |      |      |      |      |      |      |      |      |
|-----------|--|------|------|------|------|------|------|------|------|------|
| Average   | Average annual oropharyngeal cancer mortality rates* by sex, RI and US, 1987-2000**. |      |      |      |      |      |      |      |      |      |
|           | 1989   | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
| Male      |  |      |      |      |      |      |      |      |      |      |
| RI        | 7.0  | 7.1  | 6.0  | 5.4  | 5.1  | 4.9  | 4.3  | 4.3  | 4.1  | 3.5  |
| US        | 5.5  | 5.4  | 5.3  | 5.2  | 5.1  | 4.9  | 4.8  | 4.7  | 4.5  | 4.4  |
| Female    |  |      |      |      |      |      |      |      |      |      |
| RI        | 1.7  | 1.6  | 1.5  | 1.8  | 1.9  | 1.9  | 1.9  | 1.9  | 1.7  | 1.8  |
| US        | 2.0  | 2.0  | 2.0  | 1.9  | 1.9  | 1.9  | 1.8  | 1.8  | 1.7  | 1.7  |

<sup>\*</sup> Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

\*\* Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

### Table 11-5. Oropharyngeal cancer incidence by race and sex

Average annual invasive oropharyngeal cancer incidence rates\* by race and sex, RI and US, 1987-2000.

|    | M     | ale   | Female |       |  |
|----|-------|-------|--------|-------|--|
|    | White | Black | White  | Black |  |
| RI | 16.5  | 20.6  | 6.6    | 7.7   |  |
| US | 18.4  | 24.7  | 7.3    | 7.0   |  |

Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER\*Stat.

### Table 11-6. Oropharyngeal cancer mortality by race and sex

Average annual oropharyngeal cancer mortality rates\* by race and sex, RI and US, 1987-2000.

|    | M     | ale   | Female |       |  |
|----|-------|-------|--------|-------|--|
|    | White | Black | White  | Black |  |
| RI | 5.2   | 7.0   | 1.8    | 1.3   |  |
| US | 4.5   | 9.2   | 1.8    | 2.3   |  |

Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

Source: RICR, HEALTH; SEER US Mortality 1969-2000 data; calculated with SEER\*Stat.

### Tables 11-7 and 11-8. Chronic drinking by year and sex

Percent of respondents 18 and older who report an average of two or more alcoholic drinks per day (Chronic drinking = 14 or more drinks per week for males and 7 or more drinks per week for women) by year and sex, RI and US, 1990-2001.

| 1       |           |           |           |      |      | •    |      |      |      |      |      |      |
|---------|-----------|-----------|-----------|------|------|------|------|------|------|------|------|------|
|         | 1990      | 1991      | 1992      | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 |
| Male    |           |           |           |      |      |      |      |      |      |      |      |      |
| RI      | 9.5       | 9.4       | 10.2      | 8.6  | NA   | 10.0 | NA   | 7.8  | NA   | 7.3  | NA   | 8.6  |
| US      | 5.8       | 6.2       | 5.2       | 5.5  | NA   | 5.0  | NA   | 5.3  | NA   | 6.4  | NA   | 6.3  |
| Female  |           |           |           |      |      |      |      |      |      |      |      |      |
| RI      | 1.5       | 0.9       | 1.5       | 1.5  | NA   | 1.2  | NA   | 1.9  | NA   | 1.6  | NA   | 6.6  |
| US      | 0.8       | 0.9       | 8.0       | 0.9  | NA   | 8.0  | NA   | 8.0  | NA   | 0.9  | NA   | 3.9  |
| Source: | RI-BRFSS, | HEALTH; E | BRFSS, CD | C.   |      |      |      |      |      |      |      |      |

Source: Office of Vital Records, HEALTH; SEER US Mortality 1969-2000 Data; calculated with SEER\*Stat.

### Table 11-9. Oropharyngeal cancer cases in ACOS programs by year and sex

Percent of oropharyngeal cancer cases that were or are treated in ACOS-approved cancer treatment programs by year and sex, RI, 1989-2000.

|         | 1989    | 1990   | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|---------|---------|--------|------|------|------|------|------|------|------|------|------|------|
| Male    | 60.3    | 62.9   | 63.4 | 80.0 | 81.1 | 85.1 | 85.5 | 83.3 | 77.5 | 77.0 | 79.8 | 92.2 |
| Female  | 55.2    | 55.2   | 59.4 | 80.3 | 75.5 | 86.4 | 82.1 | 76.6 | 77.1 | 71.4 | 82.3 | 91.1 |
| SOURCE: | DICD HE | A I TH |      |      |      |      |      |      |      |      |      |      |

Source: RICR, HEALTH.

### Table 11-10. Oropharyngeal cancer cases with AJCC staging by year and sex

Percent of oropharyngeal cancer cases staged with AJCC staging methodology by year and sex, RI, 1989-2000.

|         | 1989      | 1990  | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|---------|-----------|-------|------|------|------|------|------|------|------|------|------|------|
| Male    | 73        | 70    | 76   | 69   | 82   | 89   | 88   | 86   | 89   | 84   | 90   | 92   |
| Female  | 48        | 61    | 73   | 71   | 78   | 83   | 86   | 90   | 78   | 78   | 76   | 85   |
| Source: | RICR, HEA | ALTH. |      |      |      |      |      |      |      |      |      |      |

### Table 11-11. Male oropharyngeal cancer survival rates by race and stage

Invasive oropharyngeal cancer five-year relative survival rates by race and stage of disease at diagnosis among males, US, 1992-1999.

|       | Localized | Regional | Distant | Unstaged |
|-------|-----------|----------|---------|----------|
| White | 82.7      | 50.3     | 25.7    | 44.1     |
| Black | 62.7      | 28.0     | 16.1    | 19.5     |

Survival rates are relative rates expressed as percents.

### Table 11-12. Female oropharyngeal cancer survival rates by race and stage

Invasive oropharyngeal cancer five-year relative survival rates by race and stage of disease at diagnosis among females, US, 1992-1999.

|       | Localized | Regional | Distant | Unstaged |
|-------|-----------|----------|---------|----------|
| White | 82.6      | 48.8     | 29.0    | 48.0     |
| Black | 77.7      | 39.1     | 34.9*   | 51.0*    |

Survival rates are relative rates expressed as percents.

### **Ovarian Cancer**

### Table 12-1. Ovarian cancer incidence by year

Average annual invasive\* ovarian cancer incidence rates\*\* among females, RI and US, 1987-2001\*\*\*.

|    | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 |
|----|------|------|------|------|------|------|------|------|------|------|------|
| RI | 16.5 | 16.3 | 16.3 | 16.5 | 17.5 | 16.6 | 16.2 | 16.1 | 16.3 | 15.4 | 14.5 |
| US | 17.5 | 17.8 | 17.8 | 17.7 | 17.6 | 17.4 | 17.3 | 17.1 | 17.2 | 16.9 | NA   |

<sup>\*</sup> Invasive includes the following stages of disease at diagnosis: local, regional, distant, and unknown.

<sup>\*\*</sup> The standard error of the survival rate is between 5 and 10 percentage points. Source: SEER Cancer Statistics Review, 1975-2000.

<sup>\*\*</sup> The standard error of the survival rate is between 5 and 10 percentage points. Source: SEER Cancer Statistics Review, 1975-2000.

<sup>\*\*</sup> Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated with SEER\*Stat.

### Table 12-2. Ovarian cancer mortality by year

Average annual ovarian cancer mortality rates\* among females, RI and US, 1987-2000\*\*.

|    | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
|----|------|------|------|------|------|------|------|------|------|------|
| RI | 9.5  | 9.2  | 8.8  | 9.2  | 8.7  | 8.3  | 8.3  | 9.0  | 8.8  | 8.8  |
| US | 9.3  | 9.4  | 9.3  | 9.4  | 9.3  | 9.2  | 9.1  | 9.0  | 8.9  | 8.8  |

Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population. Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

Office of Vital Records, HEALTH; SEER US Mortality 1969-2000 Data; calculated with SEER\*Stat.

### Table 12-3. Ovarian cancer incidence by race

Average annual invasive ovarian cancer incidence rates\* by race among females, RI and US, 1987-2000.

|    | White                       | Black                    |
|----|-----------------------------|--------------------------|
| RI | 16.4                        | 7.8                      |
| US | 18.3                        | 12.4                     |
| *  | Dates are age adjusted to t | ho year 2000 HS standard |

Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated

with SEER\*Stat.

### Table 12-4. Ovarian cancer mortality by race

Average annual ovarian cancer mortality rates\* by race among females, RI and US, 1987-2000.

|    | White                         | Black                     |
|----|-------------------------------|---------------------------|
| RI | 9.1                           | 8.4                       |
| US | 9.4                           | 7.7                       |
| *  | Datas are ago adjusted to the | ac voor 2000 IIC standard |

Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

Source: RICR, HEALTH; SEER US Mortality 1969-2000 data; calculated with SEER\*Stat.

### Table 12-5. Ovarian cancer cases in ACOS programs by year

Percent of ovarian cancer cases that were or are treated in ACOS-approved cancer treatment programs by year among females, RI, 1989-2000.

| 1989      | 1990                  | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|-----------|-----------------------|------|------|------|------|------|------|------|------|------|------|
| 48        | 42                    | 41   | 51   | 50   | 50   | 46   | 45   | 44   | 79   | 82   | 93   |
| Source: I | Source: RICR, HEALTH. |      |      |      |      |      |      |      |      |      |      |

### Table 12-6. Ovarian cancer cases with AJCC staging by year

Percent of ovarian cancer cases staged with AJCC staging methodology by year among females, RI, 1989-2000.

| 1989      | 1990                  | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |
|-----------|-----------------------|------|------|------|------|------|------|------|------|------|------|
| 73        | 68                    | 77   | 82   | 87   | 83   | 83   | 86   | 90   | 92   | 77   | 67   |
| Source: I | Source: RICR, HEALTH. |      |      |      |      |      |      |      |      |      |      |

### Table 12-7. Ovarian cancer survival rates by race and stage

Five-year relative invasive ovarian cancer survival rates by race and stage of disease at diagnosis among females, US, 1992-1999.

|  | Localized | Regional | Distant | Unstaged |  |  |  |  |  |
|--|-----------|----------|---------|----------|--|--|--|--|--|
| White  | 94.8      | 71.3     | 30.8    | 26.4     |  |  |  |  |  |
| Black  | 92.5      | 72.0*    | 26.3    | 38.4*    |  |  |  |  |  |
| * Survival rates are relative rates expressed as percents. |           |          |         |          |  |  |  |  |  |

The standard error of the survival rate is between 5 and 10

percentage points.

Source: SEER Cancer Statistics Review, 1975-2000.

### **Prostate Cancer**

### Table 13-1. Prostate cancer incidence by year

Average annual invasive\* prostate cancer incidence rates\*\* among males, RI and US, 1987-2001\*\*\*.

|    |       | •     |       |       |       |       |       |       |       |        |       |
|----|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|-------|
|    | 1989  | 1990  | 1991  | 1992  | 1993  | 1994  | 1995  | 1996  | 1997  | 1998   | 1999  |
| RI | 117.8 | 133.3 | 152.2 | 165.8 | 173.0 | 172.6 | 171.7 | 165.3 | 168.0 | 172.2  | 177.1 |
| US | 160.1 | 180.5 | 194.6 | 201.3 | 200.7 | 191.1 | 177.9 | 169.6 | 168.8 | 172.8* | NA    |

Invasive includes the following stages of disease at diagnosis: local, regional, distant, and unknown.

Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

Source: RICR, HEALTH - calculated with SEER\*Stat; SEER Cancer Statistics Review, 1973-1999;

1998 US data is from SEER Public-Use 1973-2000 Data - calculated with SEER\*Stat.

### Table 13-2. Prostate cancer mortality by year

Average annual prostate cancer mortality rates\* among males, RI and US, 1987-2000\*\*.

|    | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998  |
|----|------|------|------|------|------|------|------|------|------|-------|
| RI | 35.8 | 35.3 | 35.3 | 34.8 | 35.9 | 35.1 | 35.4 | 35.6 | 33.8 | 33.1  |
| US | 36.8 | 37.6 | 38.4 | 38.7 | 38.5 | 37.8 | 36.8 | 35.3 | 33.9 | 32.9* |

Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

Rates are based on five years' data (e.g., 1989 = 1987-1991; 1998 = 1996-2000).

Source: CDC WONDER, CDC;

1998 US data is from SEER US Mortality 1969-2000 Data - calculated with SEER\*Stat.

### Table 13-3. Prostate cancer incidence by race

Average annual invasive prostate cancer incidence rates\* by race among males, RI and US, 1987-2000.

|    | White | Black |
|----|-------|-------|
| RI | 155.9 | 206.0 |
| US | 172.3 | 267.6 |

Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Source: RICR, HEALTH; SEER Public-Use 1973-2000 Data; calculated

with SEER\*Stat.

### Table 13-4. Prostate cancer mortality by race

Average annual prostate cancer mortality rates\* by race among males, RI and US, 1987-2000.

|    | White | Black |
|----|-------|-------|
| RI | 34.3  | 77.9  |
| US | 33.2  | 75.2  |

Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

Source: RICR, HEALTH; SEER US Mortality 1969-2000 data; calculated with SEER\*Stat.

| Table 13-6   | Table 13-6. Prostate cancer cases in ACOS programs by year |      |      |      |      |      |      |      |      |      |      |  |
|--|--|------|------|------|------|------|------|------|------|------|------|--|
| Percent of prostate cancer cases that were or are treated in ACOS-approved cancer treatment programs by year among males, RI, 1989-2000. |  |      |      |      |      |      |      |      |      |      |      |  |
| 1989   | 1990   | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 |  |
| 64 56 57 78 79 78 79 80 78 78 74 88  |  |      |      |      |      |      |      |      |      |      |      |  |
| Source: RICR. HEALTH.  |  |      |      |      |      |      |      |      |      |      |      |  |

| Table 13-7  | Table 13-7. Prostate cancer cases with AJCC staging by year |  |  |  |  |  |  |  |  |  |  |  |
|---|---|--|--|--|--|--|--|--|--|--|--|--|
| Percent of prostate cancer cases staged with AJCC staging methodology by year among males, RI, 1989-2000. |   |  |  |  |  |  |  |  |  |  |  |  |
| 1989  | 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 |  |  |  |  |  |  |  |  |  |  |  |
| 51  | 51 53 52 54 66 63 73 71 69 71 73 73                         |  |  |  |  |  |  |  |  |  |  |  |
| Source: RICR, HEALTH.   |   |  |  |  |  |  |  |  |  |  |  |  |

### Table 13-8. Prostate cancer survival rates by race and stage

Five-year relative invasive prostate cancer survival rates\* by race and stage at diagnosis among males, US, 1992-1999.

| arra stag | ge at alagnosis annong                                   | , maios, 00, 1772 | .,,,,    |  |  |  |  |  |  |
|-----------|--|-------------------|----------|--|--|--|--|--|--|
|           | Local/Regional   | Distant           | Unstaged |  |  |  |  |  |  |
| White     | 100.0  | 33.3              | 89.0     |  |  |  |  |  |  |
| Black     | 100.0  | 30.4              | 85.5     |  |  |  |  |  |  |
| *         | Survival rates are relative rates expressed as percents. |                   |          |  |  |  |  |  |  |
| Source:   | SEER Cancer Statistics Review, 1975-2000.                |                   |          |  |  |  |  |  |  |

Table 14-1. Hispanic male cancer cases by data source and year

# Special Studies: Cancer Rates Among Rhode Island Hispanics

# opeolar oldares. Carroor Nates 7 arrong Nilode Island Hispanies

| Annual number of | Annual number of newly diagnosed cancers by data source and year among resident Hispanic males, RI 1989-1998. |      |      |      |      |      |      |      |      |      |  |  |
|------------------|---|------|------|------|------|------|------|------|------|------|--|--|
|                  | 1989  | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |  |  |
| Either Source    | 41  | 41   | 44   | 56   | 39   | 40   | 54   | 55   | 61   | 76   |  |  |
| Surname Analysis | 30  | 35   | 36   | 51   | 34   | 31   | 46   | 40   | 46   | 51   |  |  |
| Cancer Registry  | 21  | 16   | 20   | 10   | 14   | 17   | 18   | 28   | 36   | 44   |  |  |

\* Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population. Source: RICR, HEALTH; surname analysis technique from Word.

Table 14-2. Hispanic and all resident cancer rates by year and sex

Average annual cancer rates\* for all cancers combined by year and sex among Hispanics and all residents, RI, 1989-1998.

| Type of Rate | Sex    | Time Period | Hispanics | All Residents |
|--------------|--------|-------------|-----------|---------------|
|              |        |             | Age-Adji  | usted Rate    |
| Incidence    | Male   | 1994-1998   | 506.5     | 589.5         |
| Incidence    | Male   | 1989-1993   | 535.5     | 634.0         |
|              |        |             |           |               |
| Mortality    | Female | 1994-1998   | 143.3     | 184.4         |
| Mortality    | Female | 1989-1993   | 125.6     | 181.1         |
|              |        |             |           |               |
| Mortality    | Male   | 1994-1998   | 240.3     | 295.4         |
| Mortality    | Male   | 1989-1993   | 180.8     | 282.3         |

<sup>\*</sup> Rates are age-adjusted to the year 2000 US standard population,

expressed as cases/deaths per 100,000 population.

Source: RICR, HEALTH; Office of Vital Records, HEALTH; surname analysis technique from Word.

Table 14-3. Hispanic and all resident male cancer cases by site

Average annual percentage of all newly diagnosed cancers by anatomic site among Hispanic males and all resident males, RI, 1994-1998.

| Anatomic Site                | Hispanic Males All Resident Ma        |         |  |  |  |
|------------------------------|---------------------------------------|---------|--|--|--|
|                              | Percentage of Newly Diagnosed Cancers |         |  |  |  |
| Prostate                     | 24.8                                  | 26.9    |  |  |  |
| Lung & Bronchus              | 12.6                                  | 17.3    |  |  |  |
| Colon & Rectum               | 9.7                                   | 12.1    |  |  |  |
| Leukemias                    | 7.3                                   | 2.7     |  |  |  |
| Stomach                      | 7.0                                   | 2.4     |  |  |  |
| Lymphomas                    | 5.6                                   | 4.7     |  |  |  |
| Urinary Bladder              | 2.1                                   | 7.9     |  |  |  |
| Liver                        | 4.2                                   | 0.9     |  |  |  |
| Pancreas                     | 1.7                                   | 2.2     |  |  |  |
| Brain                        | 2.8                                   | 1.4     |  |  |  |
| Source: RICR, HEALTH; surnar | ne analysis technique fror            | n Word. |  |  |  |

# Special Studies: Cancer Mortality in Rhode Island, an Old Urban State

Cancer mortality rates\* for all cancers combined by sex and decade, RI and US, 1970-1999.

|           | 3         |           | 1         |
|-----------|-----------|-----------|-----------|
|           | 1970-1979 | 1980-1989 | 1990-1999 |
| RI Male   | 236.6     | 234.6     | 220.8     |
| US Male   | 204.3     | 212.0     | 204.0     |
| RI Female | 144.0     | 145.4     | 147.4     |
| US Female | 131.1     | 137.0     | 138.0     |

<sup>\*</sup> Rates are age-adjusted to the year 1970 US standard population, expressed as deaths per 100,000 population. Source: SEER US Mortality 1969-2000 Data; SEER web query system.

### Table 14-5. Percent elevation in white male cancer mortality by cancer site group and decade

Percent elevation in cancer mortality rates\* by cancer site group and decade among white males, RI relative to the US, 1970-1999.

|           | All | Tobacco Related | Diet Related | All Other |
|-----------|-----|-----------------|--------------|-----------|
| 1970-1979 | 16  | 18              | 28           | 3         |
| 1980-1989 | 11  | 10              | 22           | 2         |
| 1990-1999 | 8   | 10              | 14           | 2         |

\* Rates are age-adjusted to the year 1970 US standard population, expressed as deaths per 100,000 population. Source: SEER US Mortality 1969-2000 Data; SEER web query system.

### Table 14-6. Percent elevation in white female cancer mortality by cancer site group and decade

Percent elevation in cancer mortality rates\* by cancer site group and decade among white females, RI relative to the US, 1970-1999.

|           | All | Tobacco Related | Diet Related | All Other |
|-----------|-----|-----------------|--------------|-----------|
| 1970-1979 | 10  | 1               | 25           | 0         |
| 1980-1989 | 6   | 3               | 16           | 0         |
| 1990-1999 | 7   | 6               | 13           | 2         |

\* Rates are age-adjusted to the year 1970 US standard population, expressed as deaths per 100,000 population. Source: SEER US Mortality 1969-2000 Data; SEER web query system.

# Special Studies: Cancer Incidence in Rhode Island Cities and Towns

Table 14-7. Statewide and municipal cancer incidence rates by site and sex

Average annual statewide and municipal cancer incidence rates\* by anatomical site and sex, RI, 1987-2000.

|                  | -     | Ма    | ales     | -     | Females           |      |       |       |  |  |
|------------------|-------|-------|----------|-------|-------------------|------|-------|-------|--|--|
| Municipality     | Colon | Lung  | Prostate | All** | Colon Lung Breast |      |       |       |  |  |
| Rhode Island     | 82.6  | 105.8 | 153.0    | 601.4 | 56.4              | 54.3 | 130.6 | 435.7 |  |  |
|                  |       |       |          |       |                   |      |       |       |  |  |
| Barrington       | 81.1  | 87.9  | 173.3    | 581.4 | 49.8              | 42.1 | 146.4 | 436.5 |  |  |
| Bristol          | 79.3  | 101.0 | 143.5    | 576.6 | 49.2              | 36.5 | 129.7 | 375.7 |  |  |
| Burrillville     | 89.9  | 110.9 | 146.1    | 556.4 | 61.2              | 42.8 | 113.6 | 383.8 |  |  |
| Central Falls    | 83.3  | 147.5 | 117.4    | 621.4 | 55.0              | 59.0 | 101.7 | 398.4 |  |  |
| Charlestown      | 51.9  | 82.8  | 224.7    | 629.9 | 46.8              | 75.5 | 138.9 | 472.3 |  |  |
| Coventry         | 85.2  | 124.3 | 149.7    | 607.0 | 61.6              | 52.9 | 134.3 | 445.5 |  |  |
| Cranston         | 86.1  | 100.6 | 142.1    | 592.1 | 56.5              | 52.3 | 135.1 | 431.0 |  |  |
| Cumberland       | 76.9  | 81.9  | 161.1    | 545.6 | 66.0              | 42.5 | 135.2 | 429.2 |  |  |
| East Greenwich   | 90.6  | 105.8 | 240.4    | 726.1 | 68.7              | 42.3 | 175.2 | 508.5 |  |  |
| East Providence  | 82.8  | 109.1 | 146.5    | 591.7 | 62.1              | 49.8 | 127.1 | 438.6 |  |  |
| Exeter           | 51.0  | 98.0  | 125.3    | 449.0 | 52.1              | 51.5 | 120.7 | 427.8 |  |  |
| Foster           | 57.2  | 86.6  | 175.0    | 590.6 | 40.8              | 71.0 | 163.5 | 454.3 |  |  |
| Glocester        | 43.2  | 73.6  | 129.1    | 449.3 | 44.1              | 52.6 | 96.8  | 354.8 |  |  |
| Hopkinton        | 102.5 | 100.7 | 176.9    | 669.3 | 61.5              | 62.6 | 151.3 | 512.4 |  |  |
| Jamestown        | 45.8  | 80.2  | 196.3    | 623.6 | 58.9              | 66.0 | 152.4 | 476.3 |  |  |
| Johnston         | 85.4  | 113.6 | 140.0    | 604.2 | 58.5              | 56.0 | 147.1 | 468.6 |  |  |
| Lincoln          | 79.2  | 90.1  | 150.1    | 567.5 | 49.2              | 36.3 | 114.9 | 365.0 |  |  |
| Little Compton   | 65.2  | 108.1 | 169.5    | 587.2 | 29.2              | 39.7 | 141.1 | 422.0 |  |  |
| Middletown       | 70.6  | 111.7 | 236.1    | 693.8 | 58.3              | 70.2 | 146.4 | 472.2 |  |  |
| Narragansett     | 77.1  | 84.9  | 198.2    | 579.0 | 68.7              | 54.6 | 130.6 | 457.6 |  |  |
| New Shoreham     | 20.8  | 81.9  | 188.3    | 636.6 | 0.0               | 60.4 | 174.3 | 394.0 |  |  |
| Newport          | 76.4  | 107.3 | 247.6    | 716.6 | 58.3              | 72.4 | 139.9 | 480.9 |  |  |
| North Kingstown  | 96.4  | 102.2 | 207.8    | 706.0 | 50.9              | 61.9 | 139.1 | 470.7 |  |  |
| North Providence | 89.4  | 103.0 | 124.9    | 594.2 | 51.1              | 56.1 | 131.8 | 425.0 |  |  |
| North Smithfield | 72.7  | 77.5  | 118.4    | 523.7 | 64.6              | 45.0 | 115.8 | 431.0 |  |  |
| Pawtucket        | 94.5  | 113.1 | 130.8    | 605.0 | 55.7              | 52.7 | 123.2 | 428.1 |  |  |
| Portsmouth       | 87.4  | 97.1  | 202.3    | 598.5 | 63.1              | 62.4 | 153.1 | 468.0 |  |  |
| Providence       | 72.2  | 111.6 | 136.7    | 588.7 | 53.2              | 56.7 | 123.2 | 435.5 |  |  |
| Richmond         | 89.6  | 102.0 | 178.5    | 671.3 | 47.9              | 64.4 | 75.2  | 331.8 |  |  |
| Scituate         | 66.2  | 125.2 | 176.7    | 602.1 | 51.6              | 42.1 | 167.3 | 438.1 |  |  |
| Smithfield       | 74.2  | 93.9  | 151.8    | 581.3 | 54.4              | 59.5 | 130.0 | 435.3 |  |  |
| South Kingstown  | 67.9  | 99.5  | 187.0    | 608.0 | 56.0              | 57.0 | 149.4 | 447.6 |  |  |
| Tiverton         | 76.1  | 84.0  | 149.4    | 531.3 | 49.2              | 46.3 | 104.1 | 377.2 |  |  |
| Warren           | 82.1  | 98.7  | 137.5    | 575.7 | 53.4              | 59.1 | 130.6 | 427.0 |  |  |
| Warwick          | 89.8  | 114.1 | 156.8    | 637.9 | 57.8              | 65.1 | 133.7 | 460.8 |  |  |
| West Greenwich   | 91.2  | 137.3 | 182.0    | 637.6 | 46.6              | 93.6 | 109.5 | 475.5 |  |  |
| West Warwick     | 103.8 | 125.3 | 144.1    | 649.9 | 57.1              | 59.7 | 123.3 | 440.5 |  |  |
| Westerly         | 78.5  | 89.7  | 174.0    | 625.1 | 61.1              | 46.8 | 127.3 | 433.7 |  |  |
| Woonsocket       | 102.6 | 121.0 | 125.5    | 608.7 | 59.1              | 53.0 | 119.2 | 425.0 |  |  |

<sup>\*</sup> Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.

Source: RICR, HEALTH.

<sup>\*\*</sup> All cancers combined

Table 14-8. Statewide cancer incidence rates and standard deviations of municipal rates by site and sex

Average annual statewide cancer incidence rates\* and standard deviations of municipal rates by anatomical site and sex, RI, 1987-2000.

|          |                    | Males     |            | Females   |           |            |  |  |
|----------|--------------------|-----------|------------|-----------|-----------|------------|--|--|
| Site     | Statewide Standard |           | SD as # of | Statewide | Standard  | SD as # of |  |  |
|          | Rate*              | Deviation | State Rate | Rate      | Deviation | State Rate |  |  |
| Colon    | 82.6               | 17.5      | 21.1%      | 56.4      | 11.8      | 20.8%      |  |  |
| Lung     | 105.8              | 16.7      | 15.8%      | 54.3      | 11.8      | 21.8%      |  |  |
| Prostate | 153.0              | 34.6      | 22.6%      |           |           |            |  |  |
| Breast   |                    |           |            | 130.6     | 20.8      | 15.9%      |  |  |
| All**    | 601.4              | 59.2      | 9.8%       | 435.7     | 39.8      | 9.1%       |  |  |

<sup>\*</sup> Rates are age-adjusted to the year 2000 US standard population, expressed as cases per 100,000 population.
\*\* All cancers combined

Source: RICR, HEALTH.

Table 14-9. Statewide and municipal cancer cases by site and sex

Average annual statewide and municipal cases of diagnosed cancer by anatomical site and sex, RI, 1987-2000.

| Ü                | Males |      |          |       |       |      |        |       |  |
|------------------|-------|------|----------|-------|-------|------|--------|-------|--|
| Municipality     | Colon | Lung | Prostate | All*  | Colon | Lung | Breast | All*  |  |
| Rhode Island     | 5132  | 6820 | 9849     | 38346 | 5420  | 4793 | 10881  | 37981 |  |
| D - mile - t - m |       |      |          |       |       |      |        |       |  |
| Barrington       | 88    | 95   | 185      | 623   | 68    | 59   | 196    | 587   |  |
| Bristol          | 122   | 167  | 244      | 936   | 108   | 74   | 231    | 720   |  |
| Burrillville     | 75    | 94   | 120      | 473   | 80    | 49   | 128    | 454   |  |
| Central Falls    | 76    | 137  | 106      | 576   | 93    | 84   | 144    | 579   |  |
| Charlestown      | 24    | 41   | 110      | 305   | 26    | 44   | 75     | 256   |  |
| Coventry         | 151   | 232  | 274      | 1121  | 163   | 136  | 336    | 1132  |  |
| Cranston         | 472   | 561  | 814      | 3291  | 491   | 396  | 979    | 3280  |  |
| Cumberland       | 149   | 171  | 322      | 1086  | 191   | 121  | 353    | 1157  |  |
| East Greenwich   | 67    | 76   | 173      | 535   | 71    | 42   | 177    | 517   |  |
| East Providence  | 296   | 404  | 543      | 2125  | 376   | 266  | 626    | 2302  |  |
| Exeter           | 13    | 28   | 33       | 131   | 18    | 18   | 41     | 148   |  |
| Foster           | 13    | 19   | 42       | 141   | 12    | 18   | 48     | 129   |  |
| Glocester        | 20    | 41   | 56       | 217   | 23    | 29   | 61     | 208   |  |
| Hopkinton        | 37    | 42   | 66       | 262   | 31    | 31   | 75     | 255   |  |
| Jamestown        | 14    | 30   | 73       | 216   | 26    | 29   | 66     | 205   |  |
| Johnston         | 166   | 229  | 285      | 1195  | 176   | 149  | 372    | 1248  |  |
| Lincoln          | 109   | 131  | 218      | 803   | 102   | 72   | 198    | 672   |  |
| Little Compton   | 18    | 28   | 48       | 159   | 10    | 14   | 43     | 132   |  |
| Middletown       | 68    | 113  | 236      | 702   | 96    | 101  | 201    | 680   |  |
| Narragansett     | 66    | 75   | 174      | 517   | 73    | 62   | 139    | 496   |  |
| New Shoreham     | 2     | 6    | 14       | 45    | 0     | 6    | 13     | 31    |  |
| Newport          | 111   | 158  | 361      | 1064  | 141   | 161  | 285    | 1043  |  |
| North Kingstown  | 127   | 143  | 281      | 972   | 96    | 116  | 260    | 882   |  |
| North Providence | 220   | 259  | 317      | 1458  | 196   | 195  | 429    | 1440  |  |
| North Smithfield | 56    | 60   | 93       | 406   | 79    | 46   | 114    | 453   |  |
| Pawtucket        | 427   | 518  | 607      | 2760  | 406   | 353  | 761    | 2777  |  |
| Portsmouth       | 83    | 105  | 200      | 611   | 81    | 83   | 199    | 609   |  |
| Providence       | 585   | 911  | 1127     | 4860  | 717   | 668  | 1394   | 5238  |  |
| Richmond         | 23    | 30   | 47       | 177   | 14    | 17   | 27     | 107   |  |
| Scituate         | 42    | 78   | 111      | 382   | 37    | 31   | 122    | 318   |  |
| Smithfield       | 87    | 112  | 178      | 687   | 105   | 99   | 203    | 728   |  |
| South Kingstown  | 90    | 135  | 252      | 820   | 105   | 97   | 253    | 781   |  |
| Tiverton         | 82    | 89   | 158      | 563   | 64    | 61   | 133    | 481   |  |
| Warren           | 66    | 82   | 116      | 474   | 68    | 67   | 138    | 477   |  |
| Warwick          | 515   | 698  | 949      | 3788  | 518   | 549  | 1058   | 3775  |  |
| West Greenwich   | 16    | 25   | 32       | 123   | 8     | 19   | 31     | 112   |  |
| West Warwick     | 170   | 233  | 258      | 1144  | 145   | 144  | 281    | 1053  |  |
| Westerly         | 120   | 139  | 281      | 976   | 141   | 96   | 239    | 876   |  |
| Woonsocket       | 266   | 325  | 345      | 1622  | 265   | 191  | 452    | 1643  |  |
| * All cancers    |       | 020  |          |       |       |      |        |       |  |

\* All cancers combined

Source: RICR, HEALTH.

Cancer in Rhode Island

Appendix: 17-23

# Special Studies: Cancer Mortality in Rhode Island Cities and Towns

Table 14-10. Statewide and municipal cancer mortality rates with standard errors and 95% confidence limits by sex Average annual statewide and municipal cancer mortality rates\* with standard errors and 95% confidence limit by sex, RI, 1987-2000.

|                  | Males        |         |              |          |                | Females |             |              |              |             |
|------------------|--------------|---------|--------------|----------|----------------|---------|-------------|--------------|--------------|-------------|
| Municipality     | Mx Rt        | Std Err | Low CL       | Hi CL    | Sig .05        | Mx Rt   | Std Err     | Low CL       | Hi CL        | Sig .05     |
|                  | (1)          | (2)     | (3)          | (4)      | (5)            | (1)     | (2)         | (3)          | (4)          | (5)         |
| [United States]* | 268.0        | 0.1     | 267.7        | 268.3    | (NA)           | 171.7   | 0.1         | 171.5        | 171.9        | (NA)        |
| [New England]*   | 273.2        | 0.6     | 272.0        | 274.4    | (NA)           | 179.5   | 0.4         | 178.8        | 180.3        | (NA)        |
| Rhode Island     | 284.3        | 4.3     | 275.9        | 292.7    | (NA)           | 181.4   | 2.8         | 175.9        | 186.9        | (NA)        |
| Barrington       | 284.1        | 34.8    | 215.9        | 352.3    |                | 172.4   | 22.2        | 128.9        | 215.9        |             |
| Bristol          | 264.2        | 26.0    | 213.2        | 315.2    |                | 138.2   | <u>16.3</u> | <u>106.3</u> | <u>170.1</u> | Low         |
| Burrillville     | 285.4        | 36.5    | 213.9        | 356.9    |                | 153.7   | 22.6        | 109.4        | 198.0        |             |
| Central Falls    | 335.1        | 37.8    | 261.0        | 409.2    |                | 171.3   | 21.4        | 129.4        | 213.2        |             |
| Charlestown      | 239.6        | 46.4    | 148.7        | 330.5    |                | 192.0   | 36.9        | 119.7        | 264.3        |             |
| Coventry         | 290.3        | 26.6    | 238.2        | 342.4    |                | 180.3   | 16.4        | 148.2        | 212.4        |             |
| Cranston         | 278.1        | 14.3    | 250.1        | 306.1    |                | 177.5   | 9.4         | 159.1        | 195.9        |             |
| Cumberland       | 256.4        | 24.3    | 208.8        | 304.0    |                | 166.5   | 15.2        | 136.7        | 196.3        |             |
| East Greenwich   | <u>197.3</u> | 33.5    | <u>131.6</u> | 263.0    | Low            | 136.6   | 22.8        | <u>91.9</u>  | <u>181.3</u> | Low         |
| East Providence  | 291.4        | 17.9    | 256.3        | 326.5    |                | 188.8   | 11.8        | 165.7        | 211.9        | Ì           |
| Exeter           | 289.6        | 67.6    | 157.1        | 422.1    |                | 199.0   | 48.3        | 104.3        | 293.7        |             |
| Foster           | 208.7        | 64.4    | 82.5         | 334.9    |                | 181.0   | 51.0        | 81.0         | 281.0        | İ           |
| Glocester        | 235.4        | 48.2    | 140.9        | 329.9    |                | 208.1   | 38.1        | 133.4        | 282.8        |             |
| Hopkinton        | 296.8        | 56.5    | 186.1        | 407.5    |                | 186.8   | 39.5        | 109.4        | 264.2        |             |
| Jamestown        | 270.7        | 62.4    | 148.4        | 393.0    |                | 224.2   | 44.9        | 136.2        | 312.2        |             |
| Johnston         | 288.9        | 24.3    | 241.3        | 336.5    |                | 176.9   | 15.9        | 145.7        | 208.1        |             |
| Lincoln          | 267.1        | 27.6    | 213.0        | 321.2    |                | 149.2   | 17.3        | 115.3        | 183.1        |             |
| Little Compton   | 288.5        | 68.1    | 155.0        | 422.0    |                | 161.8   | 44.9        | 73.8         | 249.8        |             |
| Middletown       | 278.8        | 33.2    | 213.7        | 343.9    |                | 195.3   | 22.9        | 150.4        | 240.2        |             |
| Narragansett     | 267.7        | 37.1    | 195.0        | 340.4    |                | 193.5   | 26.2        | 142.1        | 244.9        | İ           |
| New Shoreham     | 382.0        | 183.2   | 22.9         | 741.1    |                | 246.5   | 107.7       | 35.4         | 457.6        |             |
| Newport          | 290.1        | 28.0    | 235.2        | 345.0    |                | 195.0   | 18.5        | 158.7        | 231.3        |             |
| North Kingstown  | 308.7        | 31.6    | 246.8        | 370.6    |                | 183.8   | 19.3        | 146.0        | 221.6        |             |
| North Providence | 293.2        | 22.2    | 249.7        | 336.7    |                | 182.8   | 14.4        | 154.6        | 211.0        |             |
| North Smithfield | 244.2        | 34.9    | 175.8        | 312.6    |                | 174.9   | 25.1        | 125.7        | 224.1        |             |
| Pawtucket        | 297.7        | 16.3    | 265.8        | 329.6    |                | 183.2   | 10.4        | 162.8        | 203.6        |             |
| Portsmouth       | 253.3        | 35.0    | 184.7        | 321.9    |                | 163.5   | 21.9        | 120.6        | 206.4        |             |
| Providence       | 299.7        | 12.0    | 276.2        | 323.2    |                | 192.3   | 7.8         | 177.0        | 207.6        |             |
| Richmond         | 328.1        | 87.4    | 156.8        | 499.4    |                | 225.2   | 59.4        | 108.8        | 341.6        |             |
| Scituate         | 258.8        | 43.3    | 173.9        | 343.7    |                | 174.4   | 30.5        | 114.6        | 234.2        |             |
| Smithfield       | 256.9        | 29.1    | 199.9        | 313.9    |                | 189.5   | 21.0        | 148.3        | 230.7        |             |
| South Kingstown  | 255.7        | 28.2    | 200.4        | 311.0    |                | 153.9   | 18.1        | 118.4        | 189.4        |             |
| Tiverton         | 284.6        | 34.9    | 216.2        | 353.0    |                | 160.6   | 21.6        | 118.3        | 202.9        |             |
| Warren           | 253.1        | 34.6    | 185.3        | 320.9    |                | 188.8   | 25.5        | 138.8        | 238.8        |             |
| Warwick          | 301.3        | 14.5    | 272.9        | 329.7    |                | 202.8   | <u>9.7</u>  | 183.8        | 221.8        | High        |
| West Greenwich   | 282.8        | 100.4   | 86.0         | 479.6    |                | 220.1   | 68.5        | 85.8         | 354.4        | 3           |
| West Warwick     | 306.3        | 28.0    | 251.4        | 361.2    |                | 190.3   | 17.4        | 156.2        | 224.4        |             |
| Westerly         | 249.2        | 25.5    | 199.2        | 299.2    |                | 180.9   | 18.4        | 144.8        | 217.0        |             |
| Woonsocket       | 338.2        | 22.6    | 293.9        | 382.5    | High           | 178.7   | 13.5        | 152.2        | 205.2        |             |
|                  | <u> </u>     |         | <u> </u>     | <u> </u> | ı <del>y</del> |         |             | ı            | 100 000 ma   | l<br>lotion |

<sup>(1)</sup> Rates are age-adjusted to the year 2000 US standard population, expressed as deaths per 100,000 population.

<sup>(2)</sup> Standard error of proportion for the age-adjusted rate

<sup>(3) 1.96</sup> S.E. lower confidence limit

<sup>(4) 1.96</sup> S.E. higher confidence limit

<sup>(5)</sup> Indication of significant difference from the Rhode Island state rate, P < 0.05

<sup>(</sup>NA) Not Applicable

<sup>\*</sup> Rates for US and New England from SEER US Mortality 1969-2000 Data, calculated with SEER\*Stat\* Source: Office of Vital Records, HEALTH.